




Cervical Facet Radiofrequency Neurotomy



Overview

During this minimally-invasive procedure, the physician uses heat from radio waves to treat painful facet joints in your neck. This procedure is also called radiofrequency rhizotomy. It can treat pain that doesn't respond to medications or to physical therapy.

Preparation

In preparation for the procedure, you lie on your stomach. You are given medicine to make you feel relaxed. The skin and tissue of your neck is numbed.

Inserting the Cannula

The physician inserts a tube called a "cannula." A video x-ray device called a "fluoroscope" helps guide the cannula to the medial branch nerves in your spine. These tiny nerves carry pain signals from your facet joints to your brain.

Treating the Nerves

The physician inserts an electrode through the cannula. A weak electric jolt is used to test its position. If the jolt recreates the pain but does not cause any other muscular effects, it is positioned correctly. Then the physician uses the electrode to heat the nerve. This disrupts its ability to transmit pain signals. Several nerves may be treated if necessary.

End of Procedure

When the procedure is complete, the electrode and cannula are removed. A small bandage is placed on your skin. You will be monitored for a brief time before you are allowed to go home. Your injection site may feel sore after the procedure, and you may still have neck pain. If the correct nerves were treated, you will gradually experience pain relief as you heal. This may take several weeks. Your relief may last for several months.

1. If you are taking any blood thinners, you may continue to take these prior to your procedure. There is NO need to discontinue these medications.
2. Notify staff of any allergies: iodine, anesthetic, seafood, steroids, latex or contrast dye
3. Following your procedure, apply ice for 20 minutes every 1-2 hours.
4. If you encounter soreness after your procedure and are able to take NSAIDs (Ibuprofen, Aleve, Meloxicam) you may do so. Please follow medication directions on dosage and how often you may take these.

If you have any questions following your procedure or experience any complications please, call our office.